arXiv:1606.02747v2 [math.CV] 21 Jun 2016

ANOTHER METHOD FOR PROVING A HYPERGEOMETRIC GENERATING RELATION CONTIGUOUS TO THAT OF EXTON

SHANTHA KUMARI, K.[1], **PRATHIMA, J.**[2*] **AND ARJUN K. RATHIE**[3]

ABSTRACT. The aim of this note is to establish an interesting hypergeometric generating relation contiguous to that of Exton by a short method.

2000 Mathematics subject classification : Primary 33C20 ; Secondary 33C10 , 33C15

Keywords and Phrases: Pochhammer symbol, Hypergeometric function, Contiguous function, Generating relation

1. INTRODUCTION

The generalized hypergeometric functions with p numerator and q denominator parameters is defined by [7]

$$ {}_pF_q\left[\begin{array}{ccc} \alpha_1, & \ldots, & \alpha_p \\ & & \\ \beta_1, & \ldots, & \beta_q \end{array} ; \; x \right] = {}_pF_q\left[\alpha_1, \ldots, \alpha_p; \beta_1, \ldots, \beta_q; x\right] $$

$$ = \sum_{n=0}^{\infty} \frac{(\alpha_1)_n \ldots (\alpha_p)_n}{(\beta_1)_n \ldots (\beta_q)_n} \, \frac{x^n}{n!} \tag{1} $$

where $(\alpha)_n$ denotes the Pochhammer symbol (or the raised factorial or shifted factorial, since $(1)_n = n!$) defined for every $\alpha \in \mathbb{C}$ by

$$ (\alpha)_n = \begin{cases} \alpha(\alpha+1)\ldots(\alpha+n-1), & n \in \mathbb{N} \\ 1, & n = 0 \end{cases} \tag{2} $$

For a detailed exposition of this function, we refer the standard texts of Rainville [7], Slater [8] and Exton [3].

With the help of known result [1, equ.(2)]; see also [2, p.101, equ.(5)]

$$ {}_2F_1\left[\begin{matrix} a, \; a+\frac{1}{2} \\ \frac{1}{2} \end{matrix}; x^2\right] = \frac{1}{2}\left[(1+x)^{-2a} + (1-x)^{-2a}\right] \tag{3} $$

* Corresponding Author.

Exton [4] in 1999, obtained the following interesting result.

$$\sum_{n=0}^{\infty} \frac{(d)_n \left(d+\frac{1}{2}\right)_n}{\left(\frac{1}{2}\right)_n \, n!} x^{2n} \sum_{m=0}^{n} A_m \frac{(-n)_m \left(-n+\frac{1}{2}\right)_m}{m!} y^m$$
$$= \frac{1}{2}(1+x)^{-2d} \sum_{n=0}^{\infty} A_n \frac{(d)_n \left(d+\frac{1}{2}\right)_n}{n!} \left[\frac{x^2 y}{(1+x)^2}\right]^n$$
$$+ \frac{1}{2}(1-x)^{-2d} \sum_{n=0}^{\infty} A_n \frac{(d)_n \left(d+\frac{1}{2}\right)_n}{n!} \left[\frac{x^2 y}{(1-x)^2}\right]^n \tag{4}$$

where A_m is the generalized coefficient. Also, as special case, by letting

$$A_n = \frac{(a_1)_n \ldots (a_A)_n}{(h_1)_n \ldots (h_H)_n} = \frac{((a))_n}{((h))_n}$$

he deduced the following result:

$$\sum_{0}^{\infty} \frac{(d)_n \left(d+\frac{1}{2}\right)_n}{\left(\frac{1}{2}\right)_n \, n!} x^{2n} \, {}_{A+2}F_H\left[\begin{matrix}(a),\ -n,\ -n+\frac{1}{2} \\ (h)\end{matrix}; y\right]$$
$$= \frac{1}{2}(1+x)^{-2d} \, {}_{A+2}F_H\left[\begin{matrix}(a),\ d,\ d+\frac{1}{2} \\ (h)\end{matrix}; \frac{x^2 y}{(1+x)^2}\right]$$
$$+ \frac{1}{2}(1-x)^{-2d} \, {}_{A+2}F_H\left[\begin{matrix}(a),\ d,\ d+\frac{1}{2} \\ (h)\end{matrix}; \frac{x^2 y}{(1-x)^2}\right] \tag{5}$$

In 2000, with the aid of Baileys identity [1, Equ (3)]

$${}_2F_1\left[\begin{matrix}a,\ a+\frac{1}{2} \\ \frac{3}{2}\end{matrix}; x^2\right] = \frac{1}{2x(1-2a)}\left[(1+x)^{1-2a} - (1-x)^{1-2a}\right] \tag{6}$$

Malani et al. [5] established the following interesting hypergeometric generating relation by employing the same technique used by Exton.

$$\sum_{n=0}^{\infty} \frac{(d)_n \left(d+\frac{1}{2}\right)_n}{\left(\frac{3}{2}\right)_n \, n!} x^{2n} \sum_{m=0}^{n} A_m \frac{(-n)_m \left(-n-\frac{1}{2}\right)_m}{m!} y^m$$
$$= \frac{1}{2x(1-2d)}(1+x)^{1-2d} \sum_{n=0}^{\infty} A_n \frac{(d)_n \left(d-\frac{1}{2}\right)_n}{n!} \left[\frac{x^2 y}{(1+x)^2}\right]^n$$
$$- \frac{1}{2x(1-2d)}(1-x)^{1-2d} \sum_{n=0}^{\infty} A_n \frac{(d)_n \left(d-\frac{1}{2}\right)_n}{n!} \left[\frac{x^2 y}{(1-x)^2}\right]^n \tag{7}$$

Also, as special case, by letting

$$A_n = \frac{(a_1)_n \ldots (a_A)_n}{(h_1)_n \ldots (h_H)_n} = \frac{((a))_n}{((h))_n}$$

they deduced the following result :

$$\sum_{n=0}^{\infty} \frac{(d)_n \left(d+\frac{1}{2}\right)_n}{\left(\frac{3}{2}\right)_n \, n!} \, x^{2n} \, {}_{A+2}F_H\left[\begin{matrix}(a),\ -n,\ -n-\frac{1}{2}\\ (h)\end{matrix}; y\right]$$
$$= \frac{1}{2x(1-2d)} \left\{(1+x)^{1-2d} \, {}_{A+2}F_H\left[\begin{matrix}(a),\ d,\ d-\frac{1}{2}\\ (h)\end{matrix}; \frac{x^2 y}{(1+x)^2}\right]\right.$$
$$\left. - \, (1-x)^{1-2d} \, {}_{A+2}F_H\left[\begin{matrix}(a),\ d,\ d-\frac{1}{2}\\ (h)\end{matrix}; \frac{x^2 y}{(1-x)^2}\right]\right\} \tag{8}$$

The aim of this short note is to provide another method for proving the result (7) due to Malani, without using the result (6)

2. Derivation of the result (7)

Inorder to prove the result (7), we proceed as follows:

Without loss of generality we can assume that

$$\frac{1}{2x}(1+x)^{1-2d} \sum_{n=0}^{\infty} A_n \, \frac{(d)_n \left(d-\frac{1}{2}\right)_n}{n!} \left[\frac{x^2 y}{(1+x)^2}\right]^n$$
$$- \frac{1}{2x}(1-x)^{1-2d} \sum_{n=0}^{\infty} A_n \, \frac{(d)_n \left(d-\frac{1}{2}\right)_n}{n!} \left[\frac{x^2 y}{(1-x)^2}\right]^n$$
$$= \sum_{n=0}^{\infty} a_{2n+1} \, x^{2n} \tag{9}$$

Then, it is not much difficult to see that the coefficient a_{2n+1} of x^{2n} in the expansion, after some simplification is obtained as

$$a_{2n+1} = -\sum_{m=0}^{n} A_m \, \frac{(d)_m \left(d-\frac{1}{2}\right)_m}{m!} \quad \frac{(2d+2m-1)_{2n-2m+1}}{(2n-2m+1)!} \, y^m \tag{10}$$

A simple calculation shows that

$$(2d+2m-1)_{2n-2m+1} = \frac{\Gamma(2d) \, 2^{2n} \, (d)_n \left(d+\frac{1}{2}\right)_n}{\Gamma(2d-1) \, 2^{2m} \, (d)_m \left(d-\frac{1}{2}\right)_m} \tag{11}$$

and

$$(2n-2m+1)! = \frac{2^{2n} \left(\frac{3}{2}\right)_n \, n!}{2^{2m} \, (-n)_m \left(-n-\frac{1}{2}\right)_m} \tag{12}$$

Substituting these values in (10), we get

$$a_{2n+1} = (1-2d)\,\frac{(d)_n\,\left(d+\frac{1}{2}\right)_n}{\left(\frac{3}{2}\right)_n\,n!}\quad\sum_{m=0}^{n} A_m\,\frac{(-n)_m\,\left(-n-\frac{1}{2}\right)_m}{m!}\,y^m \tag{13}$$

Finally substituting this value in (9) and dividing both sides of the resulting expression by $(1-2d)$, we get the result (7). This completes the proof of (7).

Remark : The result (7) due to Malani et al. was re-derived by Qureshi et al. [6, Equ. (2.3)], in 2002.

REFERENCES

[1] Bailey, W. N., A note on the paper by Tempest and Rosenhead. Proc. London Math. Soc., Ser 2, 51, 213-214 (1950).

[2] Erdelyi et al., Higher Transcendental Functions, Volume 1, McGraw-Hill Book Company, New York, (1953).

[3] Exton, H., Multiple Hypergeometric Integrals, Halsted Press, New York, (1976).

[4] Exton, H., A new hypergeometric generating relation, J. Indian Acad. Math., 21, 53-57, (1999).

[5] Malani, S., Rathie, A. K. and Choi. J. , Another new hypergeometric generating relation contiguous to that of Exton, Comm. Korean Math. Soc. 15(4), 691-696 (2000).

[6] Qureshi, M. I., Khan, S. and Pathan, M. A. , SOme families of Gaussian hypergeometric generating relations, Proc. 3rd Annual conference of Society for Special Functions and Their Applications, Varanasi, (India), March 4-6, (2002).

[7] Rainville,E.D., Special Functions, The Macmillan Company, Inc, New York(1960), Reprinted by Chelsea Publishing Company, New York, (1971).

[8] Slater, L.J. , Generalized Hypergeometric Functions, Cambridge University Press, Cambridge, (1966).

[1] DEPARTMENT OF MATHEMATICS, A. J. INSTITUTE OF ENGINEERING AND TECHNOLOGY, MANGALORE, KARNATAKA STATE, INDIA

E-mail address: shanthakk99@gmail.com

[2] DEPARTMENT OF MATHEMATICS, MANIPAL INSTITUTE OF TECHNOLOGY, MANIPAL, KARNATAKA STATE, INDIA

E-mail address: pamrutharaj@yahoo.co.in

[3] DEPARTMENT OF MATHEMATICS, SCHOOL OF MATHEMATICAL AND PHYSICAL SCIENCES, CENTRAL UNIVERSITY OF KERALA, TEJASWINI HILLS, PERIYE P.O., KASARAGOD, 671316, KERALA STATE, INDIA.

E-mail address: akrathie@cukerala.ac.in